

血管壁における新しい酸化ストレス関連遺伝子と
転写因子の同定

(研究課題番号 11671112)

平成 11 年度～平成 13 年度科学研究費補助金 (基盤研究 (C)(2))
研究成果報告書

平成 14 年 3 月

研究代表者 西尾 善彦
(滋賀医科大学医学部助手)



研究組織

研究代表者： 西尾 善彦 (滋賀医科大学医学部助手)
 研究分担者： なし

研究経費

平成11年度	1300	千円
平成12年度	900	千円
平成13年度	900	千円
計	3100	千円

研究発表

(1) 学会誌等

- 1: Kanazawa A, Nishio Y, Kashiwagi A, Inagaki H, Kikkawa R and Horiike K: Reduced activity of mtTFA decreases the transcription in mitochondria isolated from diabetic rat heart. *Am J Physiol Endocrinol Metab* in press
- 2: Nagai Y, Nishio Y, Nakamura T, Maegawa H, Kikkawa R and Kashiwagi A: Amelioration of high fructose-induced metabolic derangements by activation of PPAR-alpha. *Am J Physiol Endocrinol Metab* in press
- 3: Shinozaki K, Hirayama A, Nishio Y, Yoshida Y, Ohtani T, Okamura T, Masada M, Kikkawa R, Kodama K, Kashiwagi A: Coronary endothelial dysfunction in the insulin-resistant state is linked to abnormal pteridine metabolism and vascular oxidative stress. *J Am Coll Cardiol* 38:1821-1828, 2001
- 4: Suzuki E, Kashiwagi A, Nishio Y, Egawa K, Shimizu S, Maegawa H, Haneda M, Yasuda H, Morikawa S, Inubushi T, Kikkawa R: Increased arterial wall stiffness limits flow volume in the lower extremities in type 2 diabetic patients. *Diabetes Care* 24:2107-2114, 2001
- 5: Nakamura T, Kishi A, Nishio Y, Maegawa H, Egawa K, Wong NC, Kojima H, Fujimiya M, Arai R, Kashiwagi A, Kikkawa R: Insulin production in a neuroectodermal tumor that expresses islet factor-1, but not pancreatic-duodenal homeobox 1. *J Clin Endocrinol Metab* 86:1795-1800, 2001
- 6: Suzuki E, Kashiwagi A, Hidaka H, Maegawa H, Nishio Y, Kojima H, Haneda M, Yasuda H, Morikawa S, Inubushi T, Kikkawa R: H- and ³¹P-magnetic resonance spectroscopy and imaging as a new diagnostic tool to evaluate neuropathic foot ulcers in Type II diabetic patients. *Diabetologia* 43:165-172, 2000
- 7: Egawa K, Maegawa H, Shimizu S, Morino K, Nishio Y, Bryer-Ash M, Cheung AT, Kolls JK,

- Kikkawa R, Kashiwagi A: Protein-tyrosine phosphatase-1B negatively regulates insulin signaling in l6myocytes and Fao hepatoma cells. *J Biol Chem* 276:10207-10211, 2001
- 8: Hidaka H, Terada M, Maegawa H, Kojima H, Koya D, Nishio Y, Haneda M, Yasuda H, Kashiwagi A, Kikkawa R: Evaluation of a new care system provided to diabetic patients in the outpatient clinic. *Intern Med* 39:783-787, 2000
- 9: Shinozaki K, Nishio Y, Okamura T, Yoshida Y, Maegawa H, Kojima H, Masada M, Toda N, Kikkawa R, Kashiwagi A: Oral administration of tetrahydrobiopterin prevents endothelial dysfunction and vascular oxidative stress in the aortas of insulin-resistant rats. *Circ Res* 87:566-573, 2000
- 10: Suzuki E, Kashiwagi A, Nishio Y, Kojima H, Maegawa H, Haneda M, Yasuda H, Morikawa S, Inubushi T, Kikkawa R: Usefulness of waveform analysis of popliteal artery in type II diabetic patients using gated magnetic resonance 2D-cine-PC imaging and ³¹P spectroscopy. *Diabetologia* 43:1031-1038, 2000
- 11: Maeno Y, Kashiwagi A, Nishio Y, Takahara N, Kikkawa R: IDL can stimulate atherogenic gene expression in cultured human vascular endothelial cells. *Diabetes Res Clin Pract* 48:127-138, 2000
- 12: Shinozaki K, Okamura T, Nishio Y, Kashiwagi A, Kikkawa R, Toda N: Evaluation of endothelial free radical release by vascular tension responses in insulin-resistant rat aorta. *Eur J Pharmacol* 394:295-299, 2000
- 13: Kashiwagi A, Shinozaki K, Nishio Y, Maegawa H, Maeno Y, Kanazawa A, Kojima H, Haneda M, Hidaka H, Yasuda H, Kikkawa R: Endothelium-specific activation of NAD(P)H oxidase in aortas of exogenously hyperinsulinemic rats. *Am J Physiol* 277:E976-E983, 1999
- 14: Kashiwagi A, Shinozaki K, Nishio Y, Okamura T, Toda N, Kikkawa R: Free radical production in endothelial cells as a pathogenetic factor for vascular dysfunction in the insulin resistance state. *Diabetes Res Clin Pract* 45:199-203, 1999
- 15: Shinozaki K, Kashiwagi A, Nishio Y, Okamura T, Yoshida Y, Masada M, Toda N, Kikkawa R: Abnormal biopterin metabolism is a major cause of impaired endothelium-dependent relaxation through nitric oxide/O₂- imbalance in insulin-resistant rat aorta. *Diabetes* 48:2437-2445, 1999
- 16: Maegawa H, Hasegawa M, Sugai S, Obata T, Ugi S, Morino K, Egawa K, Fujita T, Sakamoto T, Nishio Y, Kojima H, Haneda M, Yasuda H, Kikkawa R, Kashiwagi A: Expression of a dominant negative SHP-2 in transgenic mice induces insulin resistance. *J Biol Chem* 274:30236-30243, 1999
- 17: Maegawa H, Shi K, Hidaka H, Iwai N, Nishio Y, Egawa K, Kojima H, Haneda M, Yasuda

H, Nakamura Y, Kinoshita M, Kikkawa R, Kashiwagi A: The 3'-untranslated region polymorphism of the gene for skeletal muscle-specific glycogen-targeting subunit of protein phosphatase 1 in the type 2 diabetic Japanese population. *Diabetes* 48:1469-1472, 1999

18: Harada N, Kashiwagi A, Nishio Y, Kikkawa R: Effects of cholesterol-lowering treatments on oxidative modification of plasma intermediate density lipoprotein plus low density lipoprotein fraction in Type 2 diabetic patients. *Diabetes Res Clin Pract* 43:111-120, 1999

19: Maegawa H, Obata T, Shibata T, Fujita T, Ugi S, Morino K, Nishio Y, Kojima H, Hidaka H, Haneda M, Yasuda H, Kikkawa R, Kashiwagi A: A new antidiabetic agent (JTT-501) rapidly stimulates glucose disposal rates by enhancing insulin signal transduction in skeletal muscle. *Diabetologia* 42:151-159, 1999

(2) 口頭発表

1: Nishio Y, Kanazawa A, Kashiwagi A, Kikkawa R: Impairment of transcription of mitochondrial DNA, but not nuclear DNA in Diabetic heart. 61st annual meeting of American Diabetes Association.

2: Shinozaki K, Kashiwagi A, Nishio Y, Yoshida Y, Masada M, Maegawa H, Okamura T, Kikkawa R: Supplement of BH4 an eNOS activator, by a gene transfer of its biosynthetic enzyme cDNA into skeletal muscle improves vascular dysfunction in insulin-resistant Rats. 61st annual meeting of American Diabetes Association.

3: Suzuki E, Kashiwagi A, Nishio Y, Maegawa H, Kikkawa R: Usefulness of waveform analysis of popliteal artery in type 2 diabetic patients using magnetic resonance 2D-cine-PC imaging. 60th annual meeting of American Diabetes Association.

4: Shinozaki K, Nishio Y, Yoshida Y, Maeno Y, Masada M, Kashiwagi A: Abnormal biopterin metabolism is crucial for inducing NO/O₂- imbalance in aortas of insulin-resistant rats. 59th annual meeting of American Diabetes Association.

5: 西尾善彦、篠崎一哉、関根理、柏木厚典 インスリン抵抗性と内皮障害 第44回日本糖尿病学会 シンポジウム

(基盤研究(C)(2))
(研究課題番号 11671112)

血管壁における新しい酸化ストレス関連遺伝子と転写因子の同定

平成十四年三月

西尾 善彦

(滋賀医科大学医学部助手)

滋賀医科大学附属図書館



2001015547