

Original Article

Association between Pulse Wave Velocity and Coronary Artery Calcification in Japanese men

the Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA)

Sayuki Torii¹, Hisatomi Arima^{1,2}, Takayoshi Ohkubo³, Akira Fujiyoshi¹, Aya Kadota^{1,2}, Naoyuki Takashima¹, Sayaka Kadowaki¹, Takashi Hisamatsu^{1,2}, Yoshino Saito¹, Naoko Miyagawa¹, Maryam Zaid¹, Yoshitaka Murakami⁴, Robert D. Abbott², Minoru Horie⁵, Katsuyuki Miura^{1,2}, Hirotugu Ueshima^{1,2}
for the SESSA Research Group

¹Department of Public Health, Shiga University of Medical Science, Otsu, Shiga, Japan

²Center for Epidemiologic Research in Asia, Shiga University of Medical Science, Otsu, Shiga, Japan

³Department of Hygiene and Public Health, Teikyo University School of Medicine, Tokyo, Japan

⁴Division of Medical Statistics, Department of Social Medicine, Toho University, Tokyo, Japan

⁵Department of Cardiovascular and Respiratory Medicine, Shiga University of Medical Science, Otsu, Shiga, Japan

Aim: Pulse wave velocity (PWV) is a simple and valid clinical method for assessing arterial stiffness. Coronary artery calcification (CAC) is an intermediate stage in the process leading to overt cardiovascular disease (CVD) and an established determinant of coronary artery disease. This study aimed to examine the association between PWV and CAC in a population-based sample of Japanese men.

Methods: This is a cross-sectional study of 986 randomly selected men aged 40-79 years from Shiga, Japan. CVD-free participants were examined from 2006 to 2008. Brachial-ankle PWV (baPWV) was measured using an automatic waveform analyzer. CAC was assessed using computed tomography. Agatston scores ≥ 10 were defined as the presence of CAC.

Results: Prevalence of CAC progressively increased with rising levels of baPWV: 20.6%, 41.7%, 56.3%, and 66.7% across baPWV quartiles <1378, 1378-1563, 1564-1849, and >1849 cm/s ($P < 0.001$ for trend). Associations remained significant after adjusting for age and other factors, including body mass index, systolic blood pressure, pulse rate, total and high-density lipoprotein cholesterol, hemoglobin A1c, drinking, smoking and exercise status, and the use of medication to treat hypertension, dyslipidemia and diabetes ($P = 0.042$ for trend). The optimal cutoff level of baPWV to detect CAC was 1612 cm/s using receiver operating characteristic curve analysis.

Conclusions: Arterial stiffness as defined by an elevated baPWV is associated with an increased prevalence of CAC in a general population-based setting among Japanese men.

J Atheroscler Thromb, 2015; 22: 1266-1277.

Key words: Arterial stiffness, Pulse wave velocity (PWV), Coronary artery calcification (CAC), Coronary artery diseases (CAD), Cardiovascular diseases (CVD)

Address for correspondence: Hisatomi Arima, Center for Epidemiologic Research in Asia, Shiga University of Medical Science, Seta Tsukinowa-cho, Otsu, Shiga, 520-2192, Japan
E-mail: arima@belle.shiga-med.ac.jp

Received: March 2, 2015

Accepted for publication: May 28, 2015

Introduction

Cardiovascular disease (CVD) is one of the leading causes of premature death in Japan and other countries¹⁻³⁾. The prevalence and incidence of CVD is expected to increase because of rapidly aging populations^{1, 3, 4)}. While current attention is largely based on

the prevention of CVD, more effective strategies could include an increased focus on the preclinical phases of CVD prior to the onset of an overt CVD event.

Coronary artery calcification (CAC), measured by computed tomography (CT), is a reliable marker of subclinical coronary atherosclerosis, which can provide morphological evidence of subclinical atherosclerotic lesions⁵⁾. Furthermore, CAC has been shown to be a strong independent predictor of future coronary events, including acute myocardial infarction, acute coronary syndrome, and other manifestations of CVD such as stroke⁶⁻⁸⁾.

Pulse wave velocity (PWV) has been shown to be a reliable measure of arterial stiffness. While carotid-femoral PWV (cfPWV) is a measure of central arterial stiffness, brachial-ankle PWV (baPWV) appears to be a combined measure of central and peripheral arterial stiffness⁹⁻¹¹⁾. Because baPWV is non-invasive, convenient to measure, and requires little technical expertise, it is widely available. Both cfPWV and baPWV have been shown to be strong predictors of clinically evident CVD events¹²⁻¹⁷⁾. However, few population-based observational studies have documented a link between either measurement and subclinical atherosclerosis, including CAC^{18, 19)}. Current evidence is largely based on studies in Western countries where the risk of coronary artery disease (CAD) is high. It is unclear if findings persist in low risk populations such as Japan. Whether more subtle relationships in low risk samples exist or can be identified is unknown.

Aim

The objective of the present study is to examine the association between baPWV and CAC in a population-based sample of Japanese men enrolled in the Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA).

Methods

Study Design and Participants

SESSA is an epidemiological study of randomly selected men from the general population of Japan. Details of the study are described elsewhere^{8, 20, 21)}. In brief, 1094 community-dwelling men aged 40-79 years underwent comprehensive physical examinations following a rigid study protocol from May, 2006 to March, 2008. After excluding participants with prior CVD ($n=99$) and those with missing information on baPWV ($n=7$), a history of stroke ($n=1$), or alcohol intake ($n=1$), a total of 986 men were included in the present cross-sectional assessment of the relationship

between baPWV and subclinical atherosclerosis. This study was approved by the Institutional Review Board of Shiga University of Medical Science (No. 17-19, 17-83), and written informed consent was obtained from all participants.

Risk Factor Measurement

Body weight and height were measured while the participants were wearing light clothing without shoes. Body mass index (BMI) was calculated as weight in kilograms divided by height in meters squared. Blood pressure and pulse rate were based on the average of two consecutive measurements in the right arm after participants emptied their bladders for urinalysis and sat quietly for at least five min using an automated sphygmomanometer (BP-8800; Omron Health Care Co. Ltd, Kyoto, Japan) with an appropriately-sized cuff. Information on medical history, medication use, the current use of cigarettes and alcohol, and exercise habits were obtained using a self-administered questionnaire followed by a review by trained research technicians. Participants were categorized as either currently or not currently drinking alcohol or smoking cigarettes. Exercise was categorized as current (≥ 1 hour/week) and not current. Estimated 10-year absolute risk of CAD was calculated using the NIPPON DATA80 Risk Chart²²⁾.

Blood Test

Venipuncture was performed early in the clinic visit following measurement of blood pressure after at least a 12-hour fast. Serum was separated by centrifugation (3000 revolutions per min, for 15 min) at 4°C within 90 min of drawing blood. Samples were sent for routine laboratory tests including the assessment of lipid profiles and glucose. Plasma glucose level was measured from NaF-treated plasma using a hexokinase glucose-6-phosphate-dehydrogenase enzymatic assay, and hemoglobin A1c (HbA1c) was measured by latex agglutination immunoassay (Kyowa Medix, Tokyo, Japan). Units of HbA1c were converted from HbA1c (JDS) to HbA1c (NGSP) using the following formula: $NGSP (\%) = 1.02 \times JDS (\%) + 0.25\%$ ²³⁾. Serum total cholesterol was measured using enzymatic assays, and high-density lipoprotein (HDL) cholesterol was measured using direct methods (Kyowa Medix, Tokyo, Japan). Serum lipids were measured at a single laboratory (Shiga Laboratory, Shiga, Japan) that has been certified for standardized lipid measurements according to the protocol of the Centers for Disease Control and Prevention/the Cholesterol Reference Method Laboratory Network (CDC/CRMLN)²⁴⁾.

Pulse Wave Velocity

baPWV was measured while the participant was in a supine position after 5 min of rest using an automatic waveform analyzer (Form I PWV/ABI; Omron Health Care Co. Ltd, Kyoto, Japan)^{16, 25)}. Pressure waveforms of the brachial and tibial arteries were recorded by an oscillometric method using occlusion and sensing cuffs adapted to both arms and ankles without clothes. The average value of left and right baPWV was used in the present analysis. Electrocardiogram was monitored with electrodes placed on both wrists. Heart sounds S1 and S2 were detected by a microphone set on the left edge of the sternum at the fourth intercostal space. The automatic device calculated the path length for baPWV by using a height-based formula, and PWV was calculated as the distance between the two arterial sites divided by the time difference between waveforms conducted from the region of origin of aorta to arms and ankles.

Coronary Artery Calcification (CAC)

CAC was measured by either electron-beam computed tomography (EBCT) in 691 sample members (70.1%) using a C-150 scanner (Imatron, South San Francisco, CA, USA) or by 16-channel multidetector row computed tomography (MDCT) in 295 sample members using an Aquilion scanner (Toshiba, Tokyo, Japan). Images were obtained from the level of the root of the aorta through the heart at a slice thickness of 3 mm with a scan time of 100 ms (EBCT) or 320 ms (MDCT). We acquired images at 70% of the cardiac cycle using electrocardiogram triggering during a single breath-hold. In this study, the amount of X-ray exposure because of CT imaging was approximately 4.7 mGly, which was almost half of that in clinical settings. Quantification of CAC was performed using a DICOM workstation and AccuImage software (AccuImage Diagnostics, South San Francisco, CA, USA). The presence of CAC was defined as a minimum of three contiguous pixels (area = 1 mm²) with density \geq 130 Hounsfield units (HU). We placed a region of interest around each high-density lesion in the epicardial coronary arteries. Peak density (HU) and area (mm²) of the individual coronary calcifications were measured. A CAC score was calculated based on the methods of Agatston²⁶⁾. An Agatston score \geq 10 is defined as definite CAC²⁷⁾. All CT images were read by a trained physician who was blinded from clinical information from the participants. The protocol described was adopted from a previous study⁸⁾ in which the reproducibility of the scans showed an intraclass correlation of 0.98²⁷⁾. Since results of the current report are similar to the results of

a stratified analysis by CT-type, the definition of CAC by EBCT and MDCT are considered equivalent. Others have also reported comparable findings from CAC assessment by EBCT and MDCT²⁸⁾.

Statistical Analysis

For the purpose of showing how study characteristics and levels of CAC vary with baPWV, data are presented by baPWV quartiles. Within quartiles, continuous study characteristics are presented as mean \pm standard deviation (SD). Dichotomous variables and the prevalence of CAC are presented as percentages. While data are presented within quartiles, tests for trend are based on modeling baPWV as an independent variable in its original continuous format. Study characteristics were modeled as dependent variables. For characteristics that are continuous, general linear regression was used. For dichotomous characteristics, including CAC prevalence, logistic regression was used. For CAC prevalence, models were adjusted for age alone and for age, BMI, systolic blood pressure, pulse rate, total and HDL cholesterol, HbA1c, current alcohol drinking (yes versus no), current smoking (yes versus no), currently exercising (yes versus no), the use of medication for the treatment of hypertension, dyslipidemia, and diabetes (yes versus no), and types of CT scanning (EBCT versus MDCT) using the forced entry method. Analyses were repeated separately for subgroups defined by age (aged 40-64 years vs 65-79 years) and use of medications for hypertension. While tests for trend were based on modeling baPWV as a continuous variable, odds ratios (OR) were also derived by comparing the odds of CAC in the top 3 quartiles of baPWV to the odds of CAC in the bottom quartile. A sensitivity analysis was also conducted after exclusion of participants with atrial fibrillation. To investigate cutoff level of baPWV to detect CAC, receiver operating characteristics (ROC) curve for the model was plotted, and the area under the ROC curve was obtained. The cutoff value of baPWV that optimizes the discriminating ability for the risk of prevalent CAC was determined as the point closest to (0, 1) on the ROC curve, which was calculated as " $(1 - \text{sensitivity})^2 + (1 - \text{specificity})^2$," or the point maximizing Youden index calculated as " $\text{sensitivity} + \text{specificity} - 1$ ".²⁹⁾ OR per 1 SD increase in 10-year absolute risk of CAD based on the NIPPON DATA80 Risk Chart for CAC was also calculated in order to compare the strength of the association with that of baPWV. All reported *P*-values are based on two-sided levels of significance. Statistical analyses were performed with SPSS Statistics 22.0 for Windows (IBM, Chicago, IL, USA).

Table 1. Averages and percentages of the study characteristics of the participants according to the quartiles of baPWV: Japanese men aged 40-79, SESSA, 2006-2008

	Quartile of baPWV (cm/s)				P value
	< 1378 246	1378-1563 247	1564-1849 247	> 1849 246	
Sample size					
Age (years)	55.5 ± 9.7	61.9 ± 9.1	66.1 ± 7.9	70.3 ± 6.3	< 0.001
BMI (kg/m ²)	23.4 ± 3.0	23.9 ± 3.2	23.5 ± 2.9	23.3 ± 2.9	0.095
Systolic blood pressure (mmHg)	120.8 ± 12.6	132.3 ± 14.3	139.8 ± 16.1	152.6 ± 17.4	< 0.001
Diastolic blood pressure (mmHg)	73.8 ± 9.1	79.9 ± 10.1	81.2 ± 9.5	84.7 ± 12.0	< 0.001
Pulse rate (bpm)	62.5 ± 8.3	62.8 ± 8.9	65.7 ± 9.7	67.8 ± 12.1	< 0.001
Total cholesterol (mg/dl)	209.0 ± 33.0	211.0 ± 33.0	210.0 ± 33.0	208.0 ± 36.0	0.792
HDL cholesterol (mg/dl)	60.7 ± 16.7	57.6 ± 16.3	59.6 ± 17.3	57.7 ± 17.3	0.112
Fasting glucose (mg/dl)	98.4 ± 17.1	99.1 ± 19.0	104.2 ± 21.6	108.3 ± 24.9	< 0.001
HbA1c (NGSP) (%)	5.78 ± 0.59	5.91 ± 0.66	6.05 ± 0.75	6.25 ± 1.03	< 0.001
Current drinkers (%)	76.8% (189)	78.5% (194)	80.6% (199)	73.2% (180)	0.150
Current smokers (%)	38.2% (94)	34.8% (86)	32.8% (81)	26.0% (64)	0.020
Currently exercising (%)	35.0% (86)	42.1% (104)	48.2% (119)	44.3% (109)	0.025
Medication for hypertension (%)	8.9% (22)	25.1% (62)	34.4% (85)	42.7% (105)	< 0.001
Medication for dyslipidemia (%)	5.3% (13)	10.5% (26)	14.2% (35)	17.9% (44)	< 0.001
Medication for diabetes (%)	2.0% (5)	6.9% (17)	10.1% (25)	17.9% (44)	< 0.001
Estimated 10-year absolute risk of CAD (%) [*]	0.16 ± 0.01	0.31 ± 0.02	0.52 ± 0.03	0.79 ± 0.04	< 0.001

Values are expressed as means ± standard deviations for continuous variables and as percentages (numbers with the characteristic) for categorical variables.

Abbreviations: baPWV, brachial-ankle pulse wave velocity; CAC, coronary artery calcification; BMI, body mass index; HDL, high-density lipoprotein; HbA1c, Hemoglobin A1c; SESSA, Shiga Epidemiological Study of Subclinical Atherosclerosis.

*Estimated using the NIPPON DATA80 Risk Chart.

Results

Table 1 shows averages and percentages of the study characteristics of the participants according to quartiles of baPWV. Participants with higher baPWV levels were older and had higher systolic blood pressure, pulse rate, fasting glucose, HbA1c, and estimated 10-year absolute risk of CAD based on the NIPPON DATA80 Risk Chart. In contrast, baPWV was inversely related to cigarette smoking and positively related to current exercising. Use of medications for the treatment of hypertension, dyslipidemia, and diabetes increased with increasing levels of baPWV. Associations with the other factors were less apparent.

The prevalence of CAC across the baPWV quartiles is given in **Table 2**. Prevalence of CAC increased consistently with increasing levels of baPWV ($p < 0.001$). Prevalence was increased more than 3-fold for men in the highest baPWV quartile versus men in the lowest quartile (66.7 versus 20.7%, $p < 0.001$). Even in the adjacent 2nd quartile, CAC prevalence was double the prevalence in the lowest quartile (41.3 versus 20.7, $p < 0.001$). The association remained significant after adjustment for age and the other confounding

factors. Relative to the lowest baPWV quartile, the age-adjusted odds of CAC increased from 1.98 (95% confidence interval: 1.30-3.00) in the 2nd quartile to 3.61 (95% confidence interval: 2.29-5.71) in the top quartile. Further adjustment for the other study features weakened the association, although CAC prevalence continued to be elevated in the top two quartiles versus the bottom quartile. In both instances, the odds of CAC was increased by at least 2-fold. In the 3rd quartile, the OR was 1.95 (95% confidence interval: 1.20-3.18). In the top quartile, it was 2.20 (95% confidence interval: 1.22-3.96). Similar results were obtained even after exclusion of participants with atrial fibrillation: multivariable ORs were 1.48, 2.01, and 2.22 in the 2nd, 3rd, and top quartiles compared with the lowest quartile (P for trend=0.031). Comparable results were also obtained after inclusion of pulse pressure as a covariate instead of systolic blood pressure into the multivariable-adjusted model: multivariable ORs were 1.53, 1.99, and 2.24 in the 2nd, 3rd, and top quartiles compared with the lowest quartile (P for trend=0.021). **Table 3** shows the association between baPWV and CAC among middle-aged and elderly men. Multivariable-adjusted ORs of CAC

Table 2. Prevalence and adjusted ORs of CAC according to quartiles of baPWV: Japanese men aged 40-79, SESSA, 2006-2008

	Quartile of baPWV (cm/s)				<i>P</i> for trend
	< 1378	1378-1563	1564-1849	> 1849	
No. of CAC/participants (CAC prevalence, %)	51/246 (20.7)	102/247 (41.3)	138/247 (55.9)	164/246 (66.7)	< 0.001
Age-adjusted OR (95% CI)	1.00	1.98 (1.30-3.00)	2.89 (1.88-4.42)	3.61 (2.29-5.71)	< 0.001
Multivariable-adjusted OR (95% CI)*	1.00	1.47 (0.94-2.30)	1.95 (1.20-3.18)	2.20 (1.22-3.96)	0.042

Abbreviations: baPWV, brachial-ankle pulse wave velocity; CAC, coronary artery calcification; CI, confidence interval; OR, odds ratio.

*Adjusted for age, BMI, systolic blood pressure, pulse rate, total cholesterol, HDL cholesterol, HbA1c, drinking, smoking, and exercise status, the use of medication to treat hypertension, dyslipidemia, or diabetes, and types of CT scanning.

Table 3. Prevalence and adjusted ORs of CAC according to quartiles of baPWV for middle-aged and elderly men: Japanese men aged 40-79, SESSA, 2006-2008

	Quartiles of baPWV (cm/s) **				<i>P</i> for trend
	< 1378	1378-1563	1564-1849	> 1849	
Middle-age (aged 40-64 years)					
No. of CAC/participants (CAC prevalence, %)	39/207 (18.8)	50/150 (33.3)	60/118 (50.8)	26/49 (53.1)	< 0.001
Multivariable-adjusted OR (95% CI)*	1.00	1.45 (0.83-2.53)	2.41 (1.27-4.54)	2.24 (0.88-5.72)	0.035
Elderly (aged 65-79 years)					
No. of CAC/participants (CAC prevalence, %)	12/39 (30.8)	52/97 (53.6)	78/129 (60.5)	138/197 (70.1)	< 0.001
Multivariable-adjusted OR (95% CI)*	1.00	1.99 (0.87-4.56)	2.56 (1.11-5.88)	3.52 (1.46-8.52)	0.032

Abbreviations: baPWV, brachial-ankle pulse wave velocity; CAC, coronary artery calcification; CI, confidence interval; OR, odds ratio.

*Adjusted for BMI, systolic blood pressure, pulse rate, total cholesterol, HDL cholesterol, HbA1c, drinking, smoking, and exercise status, the use of medication to treat hypertension, dyslipidemia, or diabetes, and types of CT scanning.

***P* values for tests of heterogeneity of the effect of baPWV on CAC between age groups were 0.670 without adjustment and 0.527 with adjustment for the other characteristics.

Table 4. Prevalence and adjusted ORs of CAC according to quartiles of baPWV among participants with and without medications to treat HT: Japanese men aged 40-79, SESSA, 2006-2008

	Quartiles of baPWV (cm/s) **				<i>P</i> for trend
	< 1378	1378-1563	1564-1849	> 1849	
Participants without medications for HT					
No. of CAC/participants (CAC prevalence, %)	39/224 (17.4)	73/185 (39.5)	82/162 (50.6)	92/141 (65.2)	< 0.001
Multivariable-adjusted OR (95% CI)*	1.00	2.16 (1.30-3.58)	2.74 (1.54-4.86)	4.01 (1.96-8.23)	0.001
Participants on medications for HT					
No. of CAC/participants (CAC prevalence, %)	12/22 (54.5)	29/62 (46.8)	56/85 (65.9)	72/105 (68.6)	0.029
Multivariable-adjusted OR (95% CI)*	1.00	0.27 (0.08-0.90)	0.46 (0.14-1.49)	0.32 (0.08-1.22)	0.155

Abbreviations: baPWV, brachial-ankle pulse wave velocity; CAC, coronary artery calcification; HT, hypertension; CI, confidence interval; OR, odds ratio.

*Adjusted for age, BMI, systolic blood pressure, pulse rate, total cholesterol, HDL cholesterol, HbA1c, drinking, smoking, and exercise status, the use of medication to treat dyslipidemia or diabetes and types of CT scanning.

***P* values for tests of heterogeneity of the effect of baPWV on CAC between age groups were 0.024 without adjustment and 0.031 with adjustment for the other characteristics.

presence increased from 1.45 (95% confidence interval: 0.83-2.53) in the 2nd quartile to 2.41 (95% confidence interval: 1.27-4.54), while slight non-significant reduction was observed in the top quartile: 2.24 (95% confidence interval: 1.27-4.54). In elderly men, linear association was observed between baPWV and CAC.

Tests for heterogeneity between the age strata with and without adjustment for the study characteristics were not significant (*P*=0.670 without adjustment and *P*=0.527 with adjustment). **Table 4** shows the association between baPWV and CAC among participants with and without medications to treat hyperten-




Fig. 1. Receiver operating characteristics (ROC) curve for brachial-ankle pulse wave velocity (baPWV) to detect coronary artery calcification (CAC)

sion. The association between baPWV and CAC was strong and continuous among participants without medications to treat hypertension while there was no clear relationship among treated hypertensives (P for heterogeneity=0.024 without adjustment and 0.031 with adjustment). **Fig. 1** shows the ROC curve to investigate the cutoff level of baPWV to detect CAC. The area under the ROC curve was 0.696 (95% confidence interval: 0.664-0.729). The optimal cutoff value of baPWV was 1612cm/s with sensitivity of 0.61 and specificity of 0.70 based both on the analysis of the distance to (0, 1) on the ROC curve (0.489) and that of the Youden index (0.314).

Crude OR for prevalent CAC per 1 SD increase in estimated 10-year absolute risk of CAD based on the NIPPON DATA80 Risk Chart was 2.28 (95% confidence interval: 1.89-2.74), while crude OR per 1 SD increase in baPWV was 2.14 (95% confidence interval: 1.95-2.65).

Discussion

This study demonstrates that baPWV is significantly associated with CAC in a general population-based setting of Japanese men. The association remains significant after adjustment for age and several risk factors commonly associated with a high risk of morbidity and mortality from CVD. To the best of our knowledge, this is the largest study to have demonstrated a link between baPWV and CAC in a general population-based sample in Japan. The present

study also confirms that associations exist in a group of individuals known to have a low risk of CAD, possibly suggesting that relationships are relevant for a broad group of individuals. To the best of our knowledge, the present analysis was also the first to conduct ROC analysis between baPWV and CAC. The optimal cutoff point of baPWV to detect CAC was approximately 1600 cm/s, which is lower than the threshold level of 1800 cm/s for prediction of future clinical CVD events in a Japanese population³⁰.

In addition to Western populations^{31, 32}, a number of hospital-based studies in Asia report a link between measures of arterial stiffness and CAC³³⁻³⁷. Limited evidence has been available from population-based studies. The Rotterdam study showed a close relationship between cfPWV and CAC among 1757 elderly individuals (with an average age of 71 years)¹⁸. Compared with participants in the lowest quartile of cfPWV, those in the highest quartile had 2.12-fold greater risk of advanced CAC before and after adjustment for confounding factors. Similarly, the Framingham Heart Study described a close link between cfPWV and calcification of the aorta and coronary artery among offspring of the original Framingham cohort and in third generation members¹⁹. In addition, cfPWV was found to be one of the strongest correlates of arterial calcification in a multivariable-adjusted model. A 1SD rise in cfPWV was associated with a 2.69-fold increase in the odds of thoracic calcification and a separate 1.47-fold increase in the odds of abdominal aortic calcification and CAC (all with $P < 0.001$). The Korean Longitudinal Study on Health and Aging also investigated the correlation between cfPWV and CAC among 1,000 elderly Korean participants (aged 65 years and older) based on age- and sex-stratified random sample³⁸. The correlation coefficient between cfPWV and CAC scores was 0.196 ($P < 0.01$). While these are studies based largely on cfPWV, they are consistent with a general association between arterial stiffness and subclinical atherosclerosis in different regions of the world where the risk of CVD can vary markedly. In a cross-sectional study of 6,009 middle-aged Korean participants of a health examination at a hospital, those in the highest quartile of baPWV had 1.63-fold greater risk of CAC after adjustment for confounding factors³⁹. ERA JUMP study also showed significant associations between baPWV and CAC prevalence among 1,131 participants from Japan, South Korea, and the United States (multivariable adjusted OR 1.36 per 1 SD rise in baPWV), but did not show separately significant associations for 235 Japanese Americans or 292 Japanese in Japan⁴⁰. The present analysis from SESSA con-

firmed the hypothesis generated from previous observational studies and demonstrated clear associations between baPWV and CAC in a general population of Japanese men.

Unfortunately, while most studies are related to cfPWV^{12-14, 41, 42)}, cfPWV measurement requires technical expertise. In contrast, measurement of baPWV is relatively simple. In Asia, it is widely available in clinical practice. Unlike cfPWV, baPWV is a less specific measure of the combination of central and peripheral arterial stiffness. In comparison, cfPWV is a measure of central arterial stiffness alone. In spite of this difference, studies have shown a close correlation between baPWV and cfPWV⁹⁻¹¹⁾. Other than subclinical disease, recent longitudinal studies have also reported that baPWV is associated with overt CVD events^{15-17, 30, 43-45)}. The present results merely extend this association to disease processes that can predate the appearance of overt CVD.

In the subgroup analyses of this study, there was no clear difference in the association of baPWV with CAC between middle-aged and elderly men (aged 40-64 years and 65-79 years). In contrast, there was significant heterogeneity in the association of baPWV with CAC according to the use of medications to treat hypertension: the association between baPWV and CAC was strong and continuous among participants without treatment while there was no clear relationship among treated hypertensives. No clear link between baPWV and CAC in treated hypertensives may be attributable to the fact that severe hypertensive subjects, who are at high-risk of CAC, are more likely to receive blood pressure lowering treatment, which may also lower baPWV levels.

While the association between baPWV and subclinical atherosclerosis is consistent with findings of associations between other measures of arterial stiffness and clinical and subclinical CVD, the current report has several limitations and raises some important questions. First, the study is cross-sectional. Because of this, it cannot be determined if high levels of baPWV predate or are the consequence of subclinical atherosclerosis. It may be that baPWV simply tracks or coincides with the development of subclinical atherosclerosis. Whether baPWV and CAC have different relationships with different forms of clinical CVD is still to be elucidated. This is relevant in Asian regions, such as Japan, where the risk of stroke is significantly greater than for CAD. Different relationships may be observed in the West where CAD is dominant. Second, the study included only men. Whether findings can be extended to Japanese women is especially interesting since the risk of CAD is

exceedingly low relative to other regions of the world. The importance of baPWV could also change as populations' age increases and as the prevalence of obesity and diabetes increases. Third, while the prognostic implication of advanced CAC is established among Caucasians in the United States, it is yet to be determined among Japanese men, as only 1 patient-based prospective study has found a predictive property for CAC among Japanese men, to our knowledge⁷⁾. In addition, assessment of CAC using MDCT requires X-ray exposure. Therefore, it might be difficult to assess CAC as a marker of subclinical coronary atherosclerosis in daily clinical practice.

Some of the cited limitations can also lead to study strengths. While only Japanese men were enrolled, heterogeneity is less than in samples composed of a variety of ethnicities and cultural preferences. The sample size is reasonably large, allowing for the assessment of associations in middle-aged and elderly men. Cases of overt CVD are excluded, which further reduces the diversity of adverse correlates that act as confounding factors in groups that are at low and high risk of morbidity and mortality from CVD. Measurement of CAC also adhered to rigid standards according to procedures in Cardiovascular Institute at University of Pittsburgh.

Conclusion

Arterial stiffness defined by high baPWV value is closely associated with CAC in a general population-based sample of Japanese men. Results suggest that baPWV can identify individuals with subclinical atherosclerosis who are also at an elevated risk of clinical vascular events and death. While measurement of baPWV is a simple, non-invasive, and low-cost method of measuring arterial stiffness, its use as an adjunct in defining CVD risk profiles warrants investigation.

Acknowledgments

Members of the SESSA Research Group

Chairperson: Hirotugu Ueshima (Center for Epidemiologic Research in Asia, Department of Public Health, Shiga University of Medical Science, Otsu, Shiga).

Co-chairpersons: Katsuyuki Miura (Department of Public Health, Shiga University of Medical Science, Otsu, Shiga).

Research members: Minoru Horie, Yasutaka Nakano, Takashi Yamamoto (Department of Cardiovascular and Respiratory Medicine, Shiga University

of Medical Science, Otsu, Shiga), Emiko Ogawa (Health Administration Center, Shiga University of Medical Science, Otsu, Shiga), Hiroshi Maegawa, Itsuko Miyazawa (Division of Endocrinology and Metabolism, Department of Medicine, Shiga University of Medical Science, Otsu, Shiga), Kiyoshi Murata (Department of Radiology, Shiga University of Medical Science, Otsu, Shiga), Kenichi Mitsunami (Shiga University of Medical Science, Otsu, Shiga), Kazuhiko Nozaki (Department of Neurosurgery, Shiga University of Medical Science, Otsu, Shiga), Akihiko Shiino (Biomedical MR Science Center, Shiga University of Medical Science, Otsu, Shiga), Isao Araki (Kusatsu Public Health Center, Kusatsu, Shiga), Teruhiko Tsuru (Department of Urology, Shiga University of Medical Science, Otsu, Shiga), Ikuo Toyama (Unit for Neuropathology and Diagnostics, Molecular Neuroscience Research Center, Shiga University of Medical Science, Otsu, Shiga), Hisakazu Ogita, Souichi Kurita (Division of Medical Biochemistry, Department of Biochemistry and Molecular Biology, Shiga University of Medical Science, Otsu, Shiga), Toshinaga Maeda (Central Research Laboratory, Shiga University of Medical Science, Otsu, Shiga), Naomi Miyamatsu (Department of Clinical Nursing Science Lecture, Shiga University of Medical Science, Otsu, Shiga), Toru Kita (Kobe City Medical Center General Hospital, Kobe, Hyogo), Takeshi Kimura (Department of Cardiovascular Medicine, Kyoto University, Kyoto), Yoshihiko Nishio (Department of Diabetes, Metabolism, and Endocrinology, Kagoshima University, Kagoshima), Yasuyuki Nakamura (Department of Living and Welfare, and Cardiovascular Epidemiology, Kyoto Women's University, Kyoto), Tomonori Okamura (Department of Preventive Medicine and Public Health, School of Medicine, Keio University, Tokyo), Akira Sekikawa, Emma JM Barinas-Mitchell (Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA, USA), Daniel Edmundowicz (Department of Medicine, Section of Cardiology, School of Medicine, Temple University, Philadelphia, PA, USA), Takayoshi Ohkubo (Department of Hygiene and Public Health, Teikyo University School of Medicine, Tokyo), Atsushi Hozawa (Preventive Medicine, Epidemiology Section, Tohoku University, Tohoku Medical Megabank Organization, Sendai, Miyagi), Nagako Okuda (Department of Health and Nutrition, University of Human Arts and Sciences, Saitama), Aya Higashiyama (Research and Development Initiative Center, National Cerebral and Cardiovascular Center, Suita, Osaka), Shinya Nagasawa (Department of Epidemiology and Public Health,

Kanazawa Medical University, Kanazawa, Ishikawa), Yoshikuni Kita (Faculty of Nursing Science, Tsuruga Nursing University, Tsuruga, Fukui), Yoshitaka Murakami (Division of Medical Statistics, Department of Social Medicine, Toho University, Tokyo), Aya Kadota (Center for Epidemiologic Research in Asia, Department of Public Health, Shiga University of Medical Science, Otsu, Shiga), Akira Fujiyoshi, Naoyuki Takashima, Takashi Kadowaki, Sayaka Kadowaki (Department of Public Health, Shiga University of Medical Science, Otsu, Shiga), Robert D. Abbott, Seiko Ohno (Center for Epidemiologic Research in Asia, Shiga University of Medical Science, Otsu, Shiga), Hisatomi Arima, Takashi Hisamatsu (Center for Epidemiologic Research in Asia, Department of Public Health, Shiga University of Medical Science, Otsu, Shiga), Naoko Miyagawa, Sayuki Torii, Yoshino Saito, Maryam Zaid, Sentaro Suzuki, and Takahiro Ito (Department of Public Health, Shiga University of Medical Science, Otsu, Shiga).

Notice of Grant Support

This study was supported by Grants-in-aid for Scientific Research (A) 13307016, (A) 17209023, (A) 21249043, (A) 23249036, (A) 25253046, (A) 15H02528, and (B) 15H04773 from the Ministry of Education, Culture, Sports, Science, and Technology, Japan, and by grant R01HL068200 from the National Institutes of Health, and by Glaxo-Smith Klein.

COI

We do not have conflicts of interest to disclose.

References

- 1) Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, Abraham J, Adair T, Aggarwal R, Ahn SY, Alvarado M, Anderson HR, Anderson LM, Andrews KG, Atkinson C, Baddour LM, Barker-Collo S, Bartels DH, Bell ML, Benjamin EJ, Bennett D, Bhalla K, Bikbov B, Bin Abdulhak A, Birbeck G, Blyth F, Bolliger I, Boufous S, Bucello C, Burch M, Burney P, Carapetis J, Chen H, Chou D, Chugh SS, Coffeng LE, Colan SD, Colquhoun S, Colson KE, Condon J, Connor MD, Cooper LT, Corriere M, Cortinovis M, de Vaccaro KC, Couser W, Cowie BC, Criqui MH, Cross M, Dabhadkar KC, Dahodwala N, De Leo D, Degenhardt L, Delossantos A, Denenberg J, Des Jarlais DC, Dharmaratne SD, Dorsey ER, Driscoll T, Duber H, Ebel B, Erwin PJ, Espindola P, Ezzati M, Feigin V, Flaxman AD, Forouzanfar MH, Fowkes FG, Franklin R, Fransen M, Freeman MK, Gabriel SE, Gakidou E, Gaspari F, Gillum RF, Gonzalez-Medina D, Halasa YA, Haring D, Harrison JE, Havmoeller R, Hay RJ,

- Hoen B, Hotez PJ, Hoy D, Jacobsen KH, James SL, Jurasaria R, Jayaraman S, Johns N, Karthikeyan G, Kassebaum N, Keren A, Khoo JP, Knowlton LM, Kobusingye O, Koranteng A, Krishnamurthi R, Lipnick M, Lipschultz SE, Ohno SL, Mabweijano J, MacIntyre MF, Mallinger L, March L, Marks GB, Marks R, Matsumori A, Matzopoulos R, Mayosi BM, McAnulty JH, McDermott MM, McGrath J, Mensah GA, Merriman TR, Michaud C, Miller M, Miller TR, Mock C, Mocumbi AO, Mokdad AA, Moran A, Mulholland K, Nair MN, Naldi L, Narayan KM, Nasseri K, Norman P, O'Donnell M, Omer SB, Ortblad K, Osborne R, Ozgediz D, Pahari B, Pandian JD, Rivero AP, Padilla RP, Perez-Ruiz F, Perico N, Phillips D, Pierce K, Pope CA, Porrini E, Pourmalek F, Raju M, Ranganathan D, Rehm JT, Rein DB, Remuzzi G, Rivara FP, Roberts T, De León FR, Rosenfeld LC, Rushton L, Sacco RL, Salomon JA, Sampson U, Sanman E, Schwebel DC, Segui-Gomez M, Shepard DS, Singh D, Singleton J, Sliwa K, Smith E, Steer A, Taylor JA, Thomas B, Tleyjeh IM, Towbin JA, Truelson T, Undurraga EA, Venketasubramanian N, Vijayakumar L, Vos T, Wagner GR, Wang M, Wang W, Watt K, Weinstock MA, Weintraub R, Wilkinson JD, Woolf AD, Wulf S, Yeh PH, Yip P, Zabetian A, Zheng ZJ, Lopez AD, Murray CJ, AlMazroa MA, Memish ZA. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: A systematic analysis for the global burden of disease study 2010. *Lancet*. 2012; 380: 2095-2128
- 2) Murray CJ, Lopez AD. Mortality by cause for eight regions of the world: Global burden of disease study. *Lancet*. 1997; 349: 1269-1276
- 3) Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, Ezzati M, Shibuya K, Salomon JA, Abdalla S, Aboyans V, Abraham J, Ackerman I, Aggarwal R, Ahn SY, Ali MK, Alvarado M, Anderson HR, Anderson LM, Andrews KG, Atkinson C, Baddour LM, Bahalim AN, Barker-Collo S, Barrero LH, Bartels DH, Basáñez MG, Baxter A, Bell ML, Benjamin EJ, Bennett D, Bernabé E, Bhalla K, Bhandari B, Bikbov B, Bin Abdulhak A, Birbeck G, Black JA, Blencowe H, Blore JD, Blyth F, Bollinger I, Bonaventure A, Boufous S, Bourne R, Boussinesq M, Braithwaite T, Brayne C, Bridgett L, Brooker S, Brooks P, Brugha TS, Bryan-Hancock C, Bucello C, Buchbinder R, Buckle G, Budke CM, Burch M, Burney P, Burstein R, Calabria B, Campbell B, Canter CE, Carabin H, Carapetis J, Carmona L, Cella C, Charlson F, Chen H, Cheng AT, Chou D, Chugh SS, Coffeng LE, Colan SD, Colquhoun S, Colson KE, Condon J, Connor MD, Cooper LT, Corriere M, Cortinovis M, de Vaccaro KC, Couser W, Cowie BC, Criqui MH, Cross M, Dabhadkar KC, Dahiya M, Dahodwala N, Damsere-Derry J, Danaei G, Davis A, De Leo D, Degenhardt L, Dellavalle R, Delossantos A, Denenberg J, Derrett S, Des Jarlais DC, Dharmaratne SD, Dherani M, Diaz-Torne C, Dolk H, Dorsey ER, Driscoll T, Duber H, Ebel B, Edmond K, Elbaz A, Ali SE, Erskine H, Erwin PJ, Espindola P, Ewoigbokhan SE, Farzadfar F, Feigin V, Felson DT, Ferrari A, Ferri CP, Fèvre EM, Finucane MM, Flaxman S, Flood L, Foreman K, Forouzanfar MH, Fowkes FG, Fransen M, Freeman MK, Gabbe BJ, Gabriel SE, Gakidou E, Ganatra HA, Garcia B, Gaspari F, Gillum RF, Gmel G, Gonzalez-Medina D, Gosselin R, Grainger R, Grant B, Groeger J, Guillemin F, Gunnell D, Gupta R, Haagsma J, Hagan H, Halasa YA, Hall W, Haring D, Haro JM, Harrison JE, Havmoeller R, Hay RJ, Higashi H, Hill C, Hoen B, Hoffman H, Hotez PJ, Hoy D, Huang JJ, Ibeaneusi SE, Jacobsen KH, James SL, Jarvis D, Jurasaria R, Jayaraman S, Johns N, Jonas JB, Karthikeyan G, Kassebaum N, Kawakami N, Keren A, Khoo JP, King CH, Knowlton LM, Kobusingye O, Koranteng A, Krishnamurthi R, Laden F, Laloo R, Laslett LL, Lathlean T, Leasher JL, Lee YY, Leigh J, Levinson D, Lim SS, Limb E, Lin JK, Lipnick M, Lipschultz SE, Liu W, Loane M, Ohno SL, Lyons R, Mabweijano J, MacIntyre MF, Malekzadeh R, Mallinger L, Manivannan S, Marques W, March L, Margolis DJ, Marks GB, Marks R, Matsumori A, Matzopoulos R, Mayosi BM, McAnulty JH, McDermott MM, McGill N, McGrath J, Medina-Mora ME, Meltzer M, Mensah GA, Merriman TR, Meyer AC, Miglioli V, Miller M, Miller TR, Mitchell PB, Mock C, Mocumbi AO, Moffitt TE, Mokdad AA, Monasta L, Montico M, Moradi-Lakeh M, Moran A, Morawska L, Mori R, Murdoch ME, Mwaniki MK, Naidoo K, Nair MN, Naldi L, Narayan KM, Nelson PK, Nelson RG, Nevitt MC, Newton CR, Nolte S, Norman P, Norman R, O'Donnell M, O'Hanlon S, Olives C, Omer SB, Ortblad K, Osborne R, Ozgediz D, Page A, Pahari B, Pandian JD, Rivero AP, Patten SB, Pearce N, Padilla RP, Perez-Ruiz F, Perico N, Pesudovs K, Phillips D, Phillips MR, Pierce K, Pion S, Polanczyk GV, Polinder S, Pope CA, Popova S, Porrini E, Pourmalek F, Prince M, Pullan RL, Ramaiah KD, Ranganathan D, Razavi H, Regan M, Rehm JT, Rein DB, Remuzzi G, Richardson K, Rivara FP, Roberts T, Robinson C, De León FR, Ronfani L, Room R, Rosenfeld LC, Rushton L, Sacco RL, Saha S, Sampson U, Sanchez-Riera L, Sanman E, Schwebel DC, Scott JG, Segui-Gomez M, Shahraz S, Shepard DS, Shin H, Shivakoti R, Singh D, Singh GM, Singh JA, Singleton J, Sleet DA, Sliwa K, Smith E, Smith JL, Stapelberg NJ, Steer A, Steiner T, Stolk WA, Stovner LJ, Sudfeld C, Syed S, Tamburlini G, Tavakkoli M, Taylor HR, Taylor JA, Taylor WJ, Thomas B, Thomson WM, Thurston GD, Tleyjeh IM, Tonelli M, Towbin JA, Truelson T, Tsilimbaris MK, Ubeda C, Undurraga EA, van der Werf MJ, van Os J, Vavilala MS, Venketasubramanian N, Wang M, Wang W, Watt K, Weatherall DJ, Weinstock MA, Weintraub R, Weisskopf MG, Weissman MM, White RA, Whiteford H, Wiebe N, Wiersma ST, Wilkinson JD, Williams HC, Williams SR, Witt E, Wolfe F, Woolf AD, Wulf S, Yeh PH, Zaidi AK, Zheng ZJ, Zonies D, Lopez AD, AlMazroa MA, Memish ZA. Disability-adjusted life years (dalys) for 291 diseases and injuries in 21 regions, 1990-2010: A systematic analysis for the global burden of disease study 2010. *Lancet*. 2012; 380: 2197-2223
- 4) Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, Amann M, Anderson HR, Andrews KG, Aryee M, Atkinson C, Bacchus LJ, Bahalim AN, Balakrishnan K, Balmes J, Barker-Collo S, Baxter A, Bell ML, Blore JD, Blyth F, Bonner C, Borges G, Bourne R, Boussinesq M, Brauer M, Brooks P, Bruce NG,

- Brunkreef B, Bryan-Hancock C, Bucello C, Buchbinder R, Bull F, Burnett RT, Byers TE, Calabria B, Carapetis J, Carnahan E, Chafe Z, Charlson F, Chen H, Chen JS, Cheng AT, Child JC, Cohen A, Colson KE, Cowie BC, Darby S, Darling S, Davis A, Degenhardt L, Dentener F, Des Jarlais DC, Devries K, Dherani M, Ding EL, Dorsey ER, Driscoll T, Edmond K, Ali SE, Engell RE, Erwin PJ, Fahimi S, Falder G, Farzadfar F, Ferrari A, Finucane MM, Flaxman S, Fowkes FG, Freedman G, Freeman MK, Gakidou E, Ghosh S, Giovannucci E, Gmel G, Graham K, Grainger R, Grant B, Gunnell D, Gutierrez HR, Hall W, Hoek HW, Hogan A, Hosgood HD, Hoy D, Hu H, Hubbell BJ, Hutchings SJ, Ibeanusi SE, Jacklyn GL, Jasrasaria R, Jonas JB, Kan H, Kanis JA, Kassebaum N, Kawakami N, Khang YH, Khatibzadeh S, Khoo JP, Kok C, Laden F, Laloo R, Lan Q, Lathlean T, Leasher JL, Leigh J, Li Y, Lin JK, Lipshultz SE, London S, Lozano R, Lu Y, Mak J, Malekzadeh R, Mallinger L, Marcenes W, March L, Marks R, Martin R, McGale P, McGrath J, Mehta S, Mensah GA, Merriman TR, Micha R, Michaud C, Mishra V, Mohd Hanafiah K, Mokdad AA, Morawska L, Mozaffarian D, Murphy T, Naghavi M, Neal B, Nelson PK, Nolla JM, Norman R, Olives C, Omer SB, Orchard J, Osborne R, Ostro B, Page A, Pandey KD, Parry CD, Passmore E, Patra J, Pearce N, Pelizzari PM, Petzold M, Phillips MR, Pope D, Pope CA, Powles J, Rao M, Razavi H, Rehfuss EA, Rehm JT, Ritz B, Rivara FP, Roberts T, Robinson C, Rodriguez-Portales JA, Romieu I, Room R, Rosenfeld LC, Roy A, Rushton L, Salomon JA, Sampson U, Sanchez-Riera L, Sanman E, Sapkota A, Seedat S, Shi P, Shield K, Shivakoti R, Singh GM, Sleet DA, Smith E, Smith KR, Stapelberg NJ, Steenland K, Stöckl H, Stovner LJ, Straif K, Straney L, Thurston GD, Tran JH, Van Dingenen R, van Donkelaar A, Veerman JL, Vijayakumar L, Weintraub R, Weissman MM, White RA, Whiteford H, Wiersma ST, Wilkinson JD, Williams HC, Williams W, Wilson N, Woolf AD, Yip P, Zielinski JM, Lopez AD, Murray CJ, Ezzati M, AlMazroa MA, Memish ZA. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: A systematic analysis for the global burden of disease study 2010. *Lancet.* 2012; 380: 2224-2260
- 5) Stolzmann P, Alkadhi H, Scheffel H, Plass A, Leschka S, Falk V, Kozerke S, Wyss C, Donati OF. Combining cardiac magnetic resonance and computed tomography coronary calcium scoring: Added value for the assessment of morphological coronary disease? *Int J Cardiovasc Imaging.* 2011; 27: 969-977
- 6) Budoff MJ, Nasir K, McClelland RL, Detrano R, Wong N, Blumenthal RS, Kondos G, Kronmal RA. Coronary calcium predicts events better with absolute calcium scores than age-sex-race/ethnicity percentiles: Mesa (multi-ethnic study of atherosclerosis). *J Am Coll Cardiol.* 2009; 53: 345-352
- 7) Yamamoto H, Ohashi N, Ishibashi K, Utsunomiya H, Kunita E, Oka T, Horiguchi J, Kihara Y. Coronary calcium score as a predictor for coronary artery disease and cardiac events in Japanese high-risk patients. *Circ J.* 2011; 75: 2424-2431
- 8) Sekikawa A, Curb JD, Edmundowicz D, Okamura T, Choo J, Fujiyoshi A, Masaki K, Miura K, Kuller LH, Shin C, Ueshima H. Coronary artery calcification by computed tomography in epidemiologic research and cardiovascular disease prevention. *J Epidemiol.* 2012; 22: 188-198
- 9) Yamashina A, Tomiyama H, Takeda K, Tsuda H, Arai T, Hirose K, Koji Y, Hori S, Yamamoto Y. Validity, reproducibility, and clinical significance of noninvasive brachial-ankle pulse wave velocity measurement. *Hypertens Res.* 2002; 25: 359-364
- 10) Sugawara J, Hayashi K, Yokoi T, Cortez-Cooper MY, DeVan AE, Anton MA, Tanaka H. Brachial-ankle pulse wave velocity: An index of central arterial stiffness? *J Hum Hypertens.* 2005; 19: 401-406
- 11) Tanaka H, Munakata M, Kawano Y, Ohishi M, Shoji T, Sugawara J, Tomiyama H, Yamashina A, Yasuda H, Sawayama T, Ozawa T. Comparison between carotid-femoral and brachial-ankle pulse wave velocity as measures of arterial stiffness. *J Hypertens.* 2009; 27: 2022-2027
- 12) Laurent S, Boutouyrie P, Asmar R, Gautier I, Laloux B, Guize L, Ducimetiere P, Benetos A. Aortic stiffness is an independent predictor of all-cause and cardiovascular mortality in hypertensive patients. *Hypertension.* 2001; 37: 1236-1241
- 13) Willum-Hansen T, Staessen JA, Torp-Pedersen C, Rasmussen S, Thijss L, Ibsen H, Jeppesen J. Prognostic value of aortic pulse wave velocity as index of arterial stiffness in the general population. *Circulation.* 2006; 113: 664-670
- 14) Vlachopoulos C, Aznaouridis K, Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness: A systematic review and meta-analysis. *J Am Coll Cardiol.* 2010; 55: 1318-1327
- 15) Ninomiya T, Kojima I, Doi Y, Fukuhara M, Hirakawa Y, Hata J, Kitazono T, Kiyohara Y. Brachial-ankle pulse wave velocity predicts the development of cardiovascular disease in a general Japanese population: The hisayama study. *J Hypertens.* 2013; 31: 477-483
- 16) Takashima N, Turin TC, Matsui K, Rumana N, Nakamura Y, Kadota A, Saito Y, Sugihara H, Morita Y, Ichikawa M, Hirose K, Kawakami K, Hamajima N, Miura K, Ueshima H, Kita Y. The relationship of brachial-ankle pulse wave velocity to future cardiovascular disease events in the general Japanese population: The takashima study. *J Hum Hypertens.* 2013; 28: 323-327
- 17) Vlachopoulos C, Aznaouridis K, Terentes-Printzios D, Ioakeimidis N, Stefanadis C. Prediction of cardiovascular events and all-cause mortality with brachial-ankle elasticity index: A systematic review and meta-analysis. *Hypertension.* 2012; 60: 556-562
- 18) van Popele NM, Mattace-Raso FU, Vliegenthart R, Grobbee DE, Asmar R, van der Kuip DA, Hofman A, de Feijter PJ, Oudkerk M, Witteman JC. Aortic stiffness is associated with atherosclerosis of the coronary arteries in older adults: The rotterdam study. *J Hypertens.* 2006; 24: 2371-2376
- 19) Tsao CW, Pencina KM, Massaro JM, Benjamin EJ, Levy D, Vasan RS, Hoffmann U, O'Donnell CJ, Mitchell GF. Cross-sectional relations of arterial stiffness, pressure pulsatility, wave reflection, and arterial calcification. *Arterio-*

- scler Thromb Vasc Biol. 2014; 34: 2495-2500
- 20) Hisamatsu T, Fujiyoshi A, Miura K, Ohkubo T, Kadota A, Kadowaki S, Kadowaki T, Yamamoto T, Miyagawa N, Zaid M, Torii S, Takashima N, Murakami Y, Okamura T, Horie M, Ueshima H, Group SR. Lipoprotein particle profiles compared with standard lipids in association with coronary artery calcification in the general Japanese population. *Atherosclerosis*. 2014; 236: 237-243
- 21) Kadota A, Miura K, Okamura T, Fujiyoshi A, Ohkubo T, Kadowaki T, Takashima N, Hisamatsu T, Nakamura Y, Kasagi F, Maegawa H, Kashiwagi A, Ueshima H, Group SR, Group NDR. Carotid intima-media thickness and plaque in apparently healthy Japanese individuals with an estimated 10-year absolute risk of CAD death according to the Japan Atherosclerosis Society (JAS) guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). *J Atheroscler Thromb*. 2013; 20: 755-766
- 22) NIPPON DATA80 Research Group. Risk assessment chart for death from cardiovascular disease based on a 19-year follow-up study of a Japanese representative population. *Circ J*. 2006; 70: 1249-1255
- 23) Kashiwagi A, Kasuga M, Araki E, Oka Y, Hanafusa T, Ito H, Tominaga M, Oikawa S, Noda M, Kawamura T, Sanke T, Namba M, Hashiramoto M, Sasahara T, Nishio Y, Kuwa K, Ueki K, Takei I, Umemoto M, Murakami M, Yamakado M, Yatomi Y, Ohashi H, Committee on the Standardization of Diabetes Mellitus Related Laboratory Testing of Japan Diabetes Society. International clinical harmonization of glycated hemoglobin in Japan: From Japan diabetes society to national glycohemoglobin standardization program values. *J Diabetes Investig*. 2012; 3: 39-40
- 24) Nakamura M, Sato S, Shimamoto T. Improvement in Japanese clinical laboratory measurements of total cholesterol and HDL-cholesterol by the US cholesterol reference method laboratory network. *J Atheroscler Thromb*. 2003; 10: 145-153
- 25) Turin TC, Kita Y, Rumana N, Takashima N, Kadota A, Matsui K, Sugihara H, Morita Y, Nakamura Y, Miura K, Ueshima H. Brachial-ankle pulse wave velocity predicts all-cause mortality in the general population: Findings from the Takashima study, Japan. *Hypertens Res*. 2010; 33: 922-925
- 26) Agatston AS, Janowitz WR, Hildner FJ, Zusmer NR, Viamonte M, Detrano R. Quantification of coronary artery calcium using ultrafast computed tomography. *J Am Coll Cardiol*. 1990; 15: 827-832
- 27) Sekikawa A, Ueshima H, Kadowaki T, El-Saied A, Okamura T, Takamiya T, Kashiwagi A, Edmundowicz D, Murata K, Sutton-Tyrrell K, Maegawa H, Evans RW, Kita Y, Kuller LH. Less subclinical atherosclerosis in Japanese men in Japan than in white men in the United States in the post-world war II birth cohort. *Am J Epidemiol*. 2007; 165: 617-624
- 28) Mao SS, Pal RS, McKay CR, Gao YG, Gopal A, Ahmadi N, Child J, Carson S, Takasu J, Sarlak B, Bechmann D, Budoff MJ. Comparison of coronary artery calcium scores between electron beam computed tomography and 64-multidetector computed tomographic scanner. *J Comput Assist Tomogr*. 2009; 33: 175-178
- 29) Perkins NJ, Schisterman EF. The inconsistency of "optimal" cutpoints obtained using two criteria based on the receiver operating characteristic curve. *Am J Epidemiol*. 2006; 163: 670-675
- 30) Munakata M, Konno S, Miura Y, Yoshinaga K, J-TOPP Study Group. Prognostic significance of the brachial-ankle pulse wave velocity in patients with essential hypertension: Final results of the J-TOPP study. *Hypertens Res*. 2012; 35: 839-842
- 31) Tillin T, Chambers J, Malik I, Coady E, Byrd S, Mayet J, Wright AR, Kooner J, Shore A, Thom S, Chaturvedi N, Hughes A. Measurement of pulse wave velocity: Site matters. *J Hypertens*. 2007; 25: 383-389
- 32) Coutinho T, Turner ST, Kullo IJ. Aortic pulse wave velocity is associated with measures of subclinical target organ damage. *JACC Cardiovasc Imaging*. 2011; 4: 754-761
- 33) Park JS, Choi UJ, Lim HS, Choi BJ, Choi SY, Yoon MH, Hwang GS, Tahk SJ, Shin JH. The relationship between coronary artery calcification as assessed by multi-detector computed tomography and arterial stiffness. *Clin Exp Hypertens*. 2011; 33: 501-505
- 34) Liu CS, Li CI, Shih CM, Lin WY, Lin CH, Lai SW, Li TC, Lin CC. Arterial stiffness measured as pulse wave velocity is highly correlated with coronary atherosclerosis in asymptomatic patients. *J Atheroscler Thromb*. 2011; 18: 652-658
- 35) Nam HJ, Jung IH, Kim J, Kim JH, Suh J, Kim HS, Kim HK, Jung YJ, Kang JW, Lee S. Association between brachial-ankle pulse wave velocity and occult coronary artery disease detected by multi-detector computed tomography. *Int J Cardiol*. 2012; 157: 227-232
- 36) Tsuchiya M, Suzuki E, Egawa K, Nishio Y, Maegawa H, Inoue S, Mitsunami K, Morikawa S, Inubushi T, Kashiwagi A. Stiffness and impaired blood flow in lower-leg arteries are associated with severity of coronary artery calcification among asymptomatic type 2 diabetic patients. *Diabetes Care*. 2004; 27: 2409-2415
- 37) Mitsutake R, Miura S, Saku K. Association between coronary artery calcification score as assessed by multi-detector row computed tomography and upstroke time of pulse wave. *Intern Med*. 2007; 46: 1833-1836
- 38) Lim S, Choi HJ, Shin H, Khang AR, Kang SM, Yoon JW, Choi SH, Jeong IK, Cho SI, Park KS, Jang HC. Subclinical atherosclerosis in a community-based elderly cohort: The Korean longitudinal study on health and aging. *Int J Cardiol*. 2012; 155: 126-133
- 39) Sung KC, Lim YH, Park S, Kang SM, Park JB, Kim BJ, Shin JH. Arterial stiffness, fatty liver and the presence of coronary artery calcium in a large population cohort. *Cardiovasc Diabetol*. 2013; 12: 162
- 40) Vishnu A, Choo J, Wilcox B, Hisamatsu T, Barinas-Mitchell EJ, Fujiyoshi A, Mackey RH, Kadota A, Ahuja V, Kadowaki T, Edmundowicz D, Miura K, Rodriguez BL, Kuller LH, Shin C, Masaki K, Ueshima H, Sekikawa A, Group EJS. Brachial-ankle pulse wave velocity is associated with coronary calcification among 1131 healthy middle-aged men. *Int J Cardiol*. 2015; 189: 67-72
- 41) Inoue N, Maeda R, Kawakami H, Shokawa T, Yamamoto H, Ito C, Sasaki H. Aortic pulse wave velocity predicts cardiovascular mortality in middle-aged and elderly Japanese. *Int J Cardiol*. 2016; 209: 10-15

- nese men. *Circ J.* 2009; 73: 549-553
- 42) Lenkey Z, Illyés M, Böcskei R, Husznai R, Sárszegi Z, Meiszterics Z, Molnár FT, Hild G, Szabados S, Cziráki A, Gaszner B. Comparison of arterial stiffness parameters in patients with coronary artery disease and diabetes mellitus using arteriograph. *Physiol Res.* 2014; 63: 429-437
- 43) Matsuoka O, Otsuka K, Murakami S, Hotta N, Yamanaka G, Kubo Y, Yamanaka T, Shinagawa M, Nunoda S, Nishimura Y, Shibata K, Saitoh H, Nishinaga M, Ishine M, Wada T, Okumiya K, Matsubayashi K, Yano S, Ichihara K, Cornélissen G, Halberg F, Ozawa T. Arterial stiffness independently predicts cardiovascular events in an elderly community -- longitudinal investiga-
tion for the longevity and aging in hokkaido county (lilac) study. *Biomed Pharmacother.* 2005; 59: S40-44
- 44) Miyano I, Nishinaga M, Takata J, Shimizu Y, Okumiya K, Matsubayashi K, Ozawa T, Sugiura T, Yasuda N, Doi Y. Association between brachial-ankle pulse wave velocity and 3-year mortality in community-dwelling older adults. *Hypertens Res.* 2010; 33: 678-682
- 45) Maeda Y, Inoguchi T, Etoh E, Kodama Y, Sasaki S, Sonoda N, Nawata H, Shimabukuro M, Takayanagi R. Brachial-ankle pulse wave velocity predicts all-cause mortality and cardiovascular events in patients with diabetes: The kyushu prevention study of atherosclerosis. *Diabetes Care.* 2014; 37: 2383-2390