Supplemental Tables for Therapeutic Drug Monitoring

Optimal Sampling Strategy and Threshold of Serum Vancomycin Concentration in Elderly Japanese Patients undergoing High-flux Hemodialysis

Akio Ogawa, BS^{a,b}; Daiki Hira, PhD^{b,c}; Masayuki Tsujimoto, PhD^d; Kohshi Nishiguchi, PhD^d; Masanori Endo, BS^a; Toshiaki Ono, BS^a; Tsuguru Hatta, MD, PhD^{a,e}; Tomohiro Terada, PhD^{b,f}; and Shin-ya Morita, PhD^b

^aDepartment of Pharmacy, Omihachiman Community Medical Center, Omihachiman, Shiga, Japan ^bDepartment of Pharmacy, Shiga University of Medical Science Hospital, Otsu, Shiga, Japan

^cCollege of Pharmaceutical Sciences, Ritsumeikan University, Kusatsu, Shiga, Japan

^dDepartment of Clinical Pharmacy, Kyoto Pharmaceutical University, Kyoto, Japan

^ePresent address: Hatta Medical Clinic, Kyoto, Japan

^fPresent address: Department of Clinical Pharmacology and Therapeutics, Kyoto University Hospital, Kyoto, Japan

Supplemental Table S1. Clinical outcomes and safety after VCM administration in each patient

Case No.	Clinical outcome of the attending physician's judgment within 30 days after the start of VCM treatment	Recurrence of MRSA infection within 60 days after discontinuation of treatment	Changes in the laboratory values after the initiation of VCM treatment
1	Improvement	-	-
2	Cure	-	-
3	Death after 14 days	-	RBC, Ht, PLT, AST, CK
4	Cure	-	-
5	Cure	-	-
6	Unchanged	-	T-Bil
7	Cure	-	-
8	Cure	-	-
9	Unchanged	Recurrence after 43 days	AST

VCM, vancomycin; RBC, red blood cell; Ht, hematocrit; PLT, platelet; AST, aspartate aminotransferase; CK, creatine kinase; T-Bil, total bilirubin

Supplemental Table S2. Correlation coefficients between AUC_{24h} and VCM dose or HD conditions

Variable		AUC ₀₋₂₄		AUC ₂₄₋₄₈	
		R	<i>P</i> -value	R	<i>P</i> -value
VCM dose (mg/kg)					
Day 1 dose (mg/kg)		0.537	0.136	0.408	0.276
Day 2 dose (mg/kg)		0.402	0.429	0.216	0.681
Day 1 + Day 2 dose (mg/kg)		0.281	0.463	0.330	0.386
Maintenance dose (mg	g/kg)	0.537	0.136	0.408	0.276
HD conditions					
Dialysis membrane	1 st HD	0.249	0.517	0.337	0.375
area (m ²)	2 nd HD	0.249	0.517	0.337	0.375
D: 1	1 st HD	-0.193	0.619	-0.136	0.727
Dialysis time (h)	2 nd HD	-0.256	0.506	-0.396	0.292
Average blood flow	1 st HD	-0.168	0.666	-0.249	0.517
rate (mL/min)	2 nd HD	0.056	0.885	0.138	0.723
Total water removal	1 st HD	-0.124	0.750	-0.203	0.600
(mL)	2 nd HD	0.142	0.716	-0.025	0.949
1.00	1 st HD	0.108	0.782	0.002	0.996
VCM removal rate (%)	2 nd HD	0.098	0.801	0.202	0.603

Pearson product-moment correlation coefficient test. VCM, vancomycin; HD, hemodialysis; AUC, area under the concentration-time curve; AUC_{0-24} , AUC calculated for 24 h after the end of the 1st HD; AUC_{24-48} , AUC calculated for 24 h before the end of the 2nd HD; R, correlation coefficient; Day 1, the first day of VCM administration; Day 2, the second day of VCM administration

Supplemental Table S3. Correlation coefficients between AUC_{24h} and serum VCM concentration at each sampling point

Variable	AUC ₀₋₂₄		$ m AUC_{24-48}$		
	R	<i>P</i> -value	R	<i>P</i> -value	
C _{pre(HD1)}	0.756	0.018	0.768	0.016	
$C_{end(HD1)}$	0.618	0.076	0.705	0.034	
C_{peak}	0.972	< 0.001	0.809	0.008	
$\mathrm{C}_{24\mathrm{h}}$	0.967	< 0.001	0.983	< 0.001	
$C_{\text{pre(HD2)}}$	0.806	0.009	0.965	< 0.001	
$C_{2h(HD2)} \\$	0.739	0.023	0.780	0.013	
$C_{end(HD2)}$	0.543	0.131	0.568	0.111	

Pearson product-moment correlation coefficient test. AUC, area under the concentration-time curve; AUC_{0-24} , AUC calculated for 24 h after the end of the 1st HD; AUC_{24-48} , AUC calculated for 24 h before the end of the 2nd HD; R, correlation coefficient